

Product

YUBINARU EC GRANULE

Ingredient and Composition

Daily dose (3 Packets) contains;

[Ingredient]	[Amount]	[Specification]
d- α -Tocopherol Acetate	300 mg	JPC
Riboflavin Butyrate	6 mg	JP
Ascorbic Acid	1000 mg	JP
Ursodeoxycholic Acid	15 mg	JP

*JP: The Japanese Pharmacopoeia; JPC: Japanese Pharmaceutical Codex

Dosage and Administration

15 years old and over: 1 pack at a time.

11 years old and over, less than 15 years old: 2/3 pack at a time.

7 years old and over, less than 11 years old: 1/2 pack at a time.

Administer in the morning and evening for twice a day dosing;
and in the morning, daytime and evening for 3 times a day dosing.

Indications

Reduces the following symptoms due to peripheral circulatory disorder:
Stiff shoulder and neck, Palsy of limb, Sensitivity to cold, Chilblain.

Reduces the following symptoms: Pigmented spot, Age spot, Pigment
deposition due to sunburn and skin irritation.

Prevention of bleeding in the following cases: Gum bleeding, Nose bleeding.

"If no improvement in the above symptoms is confirmed after one month
of use, consult a doctor, a pharmacist or a dentist."

Take vitamin EC if applicable to the followings: Physical exhaustion, De-
cline of physical strength during and after sickness, Stage of old age.

Dosage form

Yellow granules (1 Pack: 2000mg)



至誠堂製薬株式会社

SHISEIDO PHARMACEUTICAL CO., LTD.